



Synthesis and Antimicrobial Activity of Dithiocarbamates of ω -Substituted (2-naphthoxy)alkanes

SADAF ZAIDI^{1,*}, DEVDUTT CHATURVEDI², MRIDULA SAXENA¹ and RICHA SRIVASTAVA³

¹Department of Applied Chemistry, Amity School of Applied Sciences, Amity University Uttar Pradesh, Lucknow Campus, Lucknow-226028, India

²Department of Chemistry, School of Physical & Material Sciences, Mahatma Gandhi Central University, Motihari-845401, India

³Amity Institute of Pharmacy, Amity University Uttar Pradesh, Lucknow Campus, Lucknow-226028, India

*Corresponding author: E-mail: sadaf.zaidi00@gmail.com

Received: 22 March 2019;

Accepted: 30 April 2019;

Published online: 30 August 2019;

AJC-19523

A series of dithiocarbamates of ω -substituted (2-naphthoxy) alkanes was developed through condensation of 2-(2-chloro-alkoxy)-naphthalene to various kinds of aliphatic, aromatic, alicyclic, heterocyclic primary and secondary amines employing benzyl trimethyl ammonium hydroxide in catalytic quantity (Triton-B/CS₂ system) afforded desired products in high yields (82-98 %). The complete series of synthesized compounds (4-48) were evaluated for antimicrobial activity through microdilution method using various bacterial and fungal strains. The antifungal and antibacterial values were estimated as MIC values. Fluconazole and ciprofloxacin [16 to 0.03 μ g/mL] were used as the standard antifungal and antibacterial drug, respectively. Out of series of evaluated compounds, some of these compounds such as compounds 28, 29, 30, 31, 32, 33 have displayed maximum potency which is comparable to standard drugs.

Keywords: Dithiocarbamates, Antimicrobial, Amines.

INTRODUCTION


Dithiocarbamates have always received the attention of the researchers round the world because of its wide utility in areas such as pharmaceuticals [1-4], intermediate product in organic synthesis [5], for the shielding of amino groups in peptide chemistry [6,7] and as linking agents in combinatorial chemistry [8-10]. Organic dithiocarbamates have been extensively used as intermediate for the synthesis of structurally diverse synthetic intermediates/molecules of biological significance like antimalarial [11], anticholinergic [12], antimicrobial [13], antimetabolic [14], antitubercular [15], antifungal [16], anticancer [17], antioxidant [18], antiprotozoal [19], antileprosy [20], antifolates [21], antitubulin [22], antialzheimer [23], anti-HIV [24], antiproliferative [25] and anticontraceptives [26] active agents. As a useful synthon organic dithiocarbamates have been extensively used for the synthesis of structurally diverse biological potent synthetic intermediates/molecules like isothiocyanates [27], thiourea [28], cyanamide [29], dithio-benzophene [30], glycosides [31], β -sulphonamides [32], amide [33], dicarboxylates [34], thiazidizoles [35], dithiolanes [36],

thiones [37], benzimidazole [38], carbamate [39], pyran [40] and flavonoids [41], *etc.* Apart from above mentioned activities, dithiocarbamates of various imidazole [42], brassinin [43], rhodanine [44], quinoline [45], metal complexes [46], ammonium salts [47], *etc.* derivative have emerged as potent antimicrobial agents. As our group is working in drug discovery through design and synthesis of novel class of natural/semi-synthetic/synthetic molecules especially molecules like carbamates, dithiocarbamates, dithiocarbazates, *etc.* Keeping in view the importance of dithiocarbamates and its derivatives, we became interested in investigating various structurally diverse biologically potent compounds.

Considering the potency of dithiocarbamates as antimicrobial agents, we became interested to investigate the antimicrobial activity of dithiocarbamates of ω -substituted (2-naphthoxy) alkanes (**Prototype I**).

EXPERIMENTAL

Procedure for ω -substituted 2-naphthoxy haloalkanes (3): Measured amount of β -naphthol (1) was taken in dry acetone



and anhyd. K_2CO_3 (10 eqv.) was added in it. To this, 1-bromo-3-chloro propane (**2**) was added (2.5 eqv.) and then the reaction was refluxed for 12-15 h. The continuous monitoring of progress of reaction was done by TLC which indicates the appearance of less polar new spot. The filtrate of reaction mixture was concentrated, extracted thrice with ethyl acetate. After separation the organic layer was dried over anhydrous Na_2SO_4 afforded the corresponding compound **3**. Compound **3** was confirmed by various spectroscopic and analytical techniques.

Procedure for synthesis of dithiocarbamates of Prototype I (4-48): Measured amount of desired amine was dissolved in dry DMSO. To this, measured amount of CS_2 and Triton-B were added drop by drop and the reaction was allowed to stir for 15 min. After adding compound **3** to the reaction mixture and the reaction was stirred for about 20-40 min. Monitoring of reaction progress was done by TLC. Triple extraction of reaction mixture was done with ethyl acetate once the reaction was complete. The organic layer was separated, dried over anhydrous Na_2SO_4 afforded the final product that is, dithiocarbamates of prototype **I** (compounds **4-48**).

Biological activities: The antibacterial and antifungal activities of the ω -substituted (2-naphthyl) alkanes against bacterial strains (*Staphylococcus aureus* ATCC 29313, Methicillin resistant *Staphylococcus aureus*), two Gram-negative strains (*Escherichia coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853), two yeast strains (*Candida albicans* ATCC 22019, *S. schenckii*) and two filamentous fungi (*Aspergillus fumigatus* LSI-II, *Aspergillus niger* ATCC 16404) was performed by using microdilution method. Antibacterial tests were performed using Muller Hinton Broth which was buffered to pH 7.0. The antifungal testing was performed using RPMI 1640 with L-glutamine buffered to pH 7.0. It was supplemented with 0.165 M 3-(*N*-morpholino) propanesulfonic acid (MOPS) [Sigma-Aldrich]. The stock solution of the compounds was prepared using DMSO. The minimum inhibitory concentration (MIC) of the compounds was determined by serial 2 fold diluting the oils in the abovementioned media in 100 mL volume in a 96 well U bottom microtitre plate. The final concentrations of compound ranged from 128 to 0.25 $\mu g/mL$. Fluconazole and ciprofloxacin [16 to 0.03 $\mu g/mL$] (Sigma-Aldrich) were used as standard antifungal and antibacterial agents respectively. The bacterial and fungal suspension of the overnight grown bacterial and fungal was prepared in sterile normal

saline and their density was adjusted to 0.5 McFarland. The bacterial cultures were diluted and added in 100 mL volume to a final inoculum of 1×10^5 CFU/mL. For fungal cultures 1×10^3 CFU/mL was used. The plates were incubated at 37 $^\circ C$ for 24 h for bacterial cultures and 48 h for fungal cultures. The plates were read visually and the minimum concentration of the compounds showing no turbidity was recorded as MIC.

All the chemicals used were obtained from Merck, Aldrich and Fluka chemical companies. Reactions were carried out in nitrogenous atmosphere. The structural analysis of compound was done as IR spectra (4000-200 cm^{-1}) on Bomem MB-104-FTIR spectrophotometer where as NMRs were scanned on AC-300F, NMR (300 MHz), instrument using $CDCl_3$ and some other deuterated solvents and TMS as internal standard. Elemental analysis were made by Carlo-Erba EA 1110-CNNO-S analyzer and the obtained values were in accordance with calculated values.


General procedure for the preparation of ω -naphthyl halo alkanes (**3a-c**) involves refluxing of mixture of β -naphthol (**1**) (20 g, 0.14 mol), anhydrous K_2CO_3 (100 g, in excess) and bromochloroalkane **2** (0.14 mol) in dry acetone (200 mL) for 12-15 h. Reaction mixture was filtered and filtrate was concentrated to get oily compound, which was crystallized with benzene-hexane to give the colourless crystals of pure desired compound (**Scheme-I**) [48].

2-(2-Naphthyl)oxy-1-chloroethane (3a): Yield: 27.4 g (96 %); m.p.: 94 $^\circ C$; IR (KBr, ν_{max} , cm^{-1}): 1455 (Ar), 1507 (Ar), 1586 (Ar), 2878 (CH), 2927 (CH); 1H NMR (400 MHz, $CDCl_3$): $\delta = 3.82$ (t, 2H, CH_2Cl), 4.24 (t, 2H, OCH_2), 6.97–7.65 (m, 7H, Ar-H); ^{13}C NMR (100 MHz, $CDCl_3$): $\delta = 45.2, 75.2, 105.7, 118.5, 123.7, 126.5, 129.6, 134.4, 157.6$ ppm; Mass (EIMS): $m/z = 206$; Analysis: $C_{12}H_{11}OCl$, Calcd. (%): C, 69.74; H, 5.36; Obsd. (%): C, 70.04, H, 5.66.

3-(2-Naphthyl)oxy-1-chloropropane (3b): Yield: 29.6 g (97 %); m.p.: 98 $^\circ C$; IR (KBr, ν_{max} , cm^{-1}): 1462 (Ar), 1511 (Ar), 1595 (Ar), 2856 (CH), 2942 (CH); 1H NMR (400 MHz, $CDCl_3$): $\delta = 2.26$ –2.34 (m, 2H, CH_2), 3.84 (t, 2H, CH_2Cl), 4.24 (t, 2H, OCH_2), 7.12–7.75 (m, 7H, Ar-H) ppm; Mass (EIMS): $m/z = 220$; Analysis: $C_{13}H_{13}OCl$, Calcd. (%): C, 70.75; H, 5.94; Obsd. (%): C, 70.79; H, 6.21.

4-(2-Naphthyl)oxy-1-chlorobutane (3c): Yield: 34 g (98 %); m.p.: 112 $^\circ C$; IR (KBr, ν_{max} , cm^{-1}): 1463 (Ar), 1512 (Ar), 1598 (Ar), 2886 (CH), 2942 (CH); 1H NMR (400 MHz, $CDCl_3$): $\delta = 2.15$ –2.22 (m, 4H, CH_2CH_2), 3.77 (t, 2H, CH_2Cl), 4.25 (t, 2H, OCH_2), 7.13–7.76 (m, 7H, Ar-H) ppm; Mass (EIMS): $m/z = 234$; Analysis: $C_{14}H_{15}OCl$, Calcd. (%): C, 70.72; H, 5.92; Obsd. (%): C, 70.78; H, 6.20.

Procedure for the preparation of dithiocarbamates of ω -substituted (2-naphthyl) alkanes: A mixture of desired



Scheme-I

amine (0.6 mL, 5 mmol) and carbon disulphide solution (3 mL, in excess) was taken in dry DMSO (35 mL). Triton-B (0.9 mL, 4 mmol) was added in it and the reaction mixture was stirred at room temperature for 1 h. Now 2-(2-naphthylloxy)-1-chloroalkane (0.5 g, 2 mmol) was added in it and reaction was continued till its completion (2 h) as checked by TLC. Reaction mixture was poured into distilled water (50 mL) and three-time extraction was done with ethyl acetate. After separation of organic layer, it was dried over anhydrous sodium sulphate and then concentrated to get dithiocarbamate of ω -substituted (2-naphthylloxy) alkanes. This compound was obtained as yellow solid (Scheme-II).

Butyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (4): Yield: 0.73 g (93.5 %); m.p.: 106 °C; IR (KBr, ν_{\max} , cm^{-1}): 661 (C-S), 1114 (C=S), 1454 (Ar), 1511 (Ar), 1612 (Ar), 2864 (CH), 2936 (CH), 3390 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.96 (t, 3H, CH_3), 1.30-1.34 (m, 2H, CH_2CH_3), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 2.0 (bs, H, NH), 2.62-2.64 (m, 2H, NHCH_2), 3.28-3.32 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.71-4.74 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.97-7.64 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 319; Analysis: $\text{C}_{17}\text{H}_{21}\text{NOS}_2$, Calcd. (%): C, 63.91, H 6.63, N, 4.38; Obsd. (%): C, 64.19, H, 6.49, N, 4.24.

Butyl-dithiocarbamic acid-3-(naphthalene-2-yloxy)-propyl ester (5): Yield: 0.73 g (93.5 %); m.p.: 106 °C; IR (KBr, ν_{\max} , cm^{-1}): 661 (C-S), 1115 (C=S), 1454 (Ar), 1511 (Ar), 1610 (Ar), 2864 (CH), 2935 (CH), 3390 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.93-0.96 (t, 3H, CH_3), 1.33-1.35 (m, 2H, CH_2CH_3), 1.53-1.55 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 2.0 (bs, H, NH), 2.62-2.65 (m, 2H, NHCH_2), 2.84-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 2.35-2.39 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.01-4.05 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 333; Analysis: $\text{C}_{18}\text{H}_{23}\text{NOS}_2$, Calcd. (%): C, 65.91, H 6.83, N, 4.38, S, 19.15; Obsd. (%): C, 65.82, H, 6.95, N, 4.20, S, 19.23 %; O, 4.80.

Butyl-dithiocarbamic acid-4-(naphthalene-2-yloxy)-butyl ester (6): Yield: 0.73 g (93.5 %); m.p.: 106 °C; IR (KBr, ν_{\max} , cm^{-1}): 661 (C-S), 1114 (C=S), 1454 (Ar), 1511 (Ar), 1612 (Ar), 2864 (CH), 2936 (CH), 3390 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.93-0.96 (t, 3H, CH_3), 1.33-1.35 (m, 2H, CH_2CH_3), 1.53-1.55 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$), 2.0 (bs, H, NH), 2.62-2.66 (m, 2H, NHCH_2), 2.85-2.87 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.92-1.96 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.68-1.71 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.00-4.03 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 347.14; Analysis: $\text{C}_{19}\text{H}_{25}\text{NOS}_2$, Calcd. (%): C, 65.60, H 7.22, N, 4.58; S, 18.42 %; O, 4.58. Obsd. (%): C, 65.66, H, 7.25, N, 4.60, S, 18.45 %; O, 4.60.


Hexyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (7): Yield: 0.8 g (96.4 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm^{-1}): 665 (C-S), 1116 (C=S), 1475 (Ar), 1514 (Ar), 1602

(Ar), 2875 (CH), 2936 (CH), 3394 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.95 (t, 3H, CH_3), 1.27-1.29 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 1.31-1.35 (m, 2H, CH_2CH_3 of hexyl group), 1.52-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 2.0 (bs, H, NH), 2.36-2.40 (m, 2H, naphthyl-O- $\text{CH}_2\text{CH}_2\text{CH}_2\text{-S-C=S}$), 2.63-2.65 (m, 2H, NHCH_2), 3.24-3.29 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.68-4.73 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 347.14; Analysis: $\text{C}_{19}\text{H}_{25}\text{NOS}_2$, Calcd. (%): C, 66.44, H 7.53, N, 3.87, 18.45 %; O, 4.60. Obsd. (%): C, 65.66, H, 7.25, N, 4.03, S, 18.45 %; O, 4.60.

Hexyl-dithiocarbamic acid-3-(naphthalene-2-yloxy)-propyl ester (8): Yield: 0.8 g (96.4 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1116 (C=S), 1475 (Ar), 1514 (Ar), 1602 (Ar), 2875 (CH), 2936 (CH), 3394 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.94 (t, 3H, CH_3), 1.27-1.29 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 1.31-1.35 (m, 2H, CH_2CH_3 of hexyl group), 1.52-1.58 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 2.0 (bs, H, NH), 2.34-2.42 (m, 2H, naphthyl-O- $\text{CH}_2\text{CH}_2\text{CH}_2\text{-S-C=S}$), 2.63-2.66 (m, 2H, NHCH_2), 2.81-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.02-4.05 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.97-7.64 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 361; Analysis: $\text{C}_{20}\text{H}_{27}\text{NOS}_2$, Calcd. (%): C, 66.44, H 7.53, N, 3.87; Obsd. (%): C, 66.75, H, 7.38, N, 3.71.

Hexyl-dithiocarbamic acid-4-(naphthalene-2-yloxy)-butyl ester (9): Yield: 0.72 g (98 %); m.p.: 129 °C; IR (KBr, ν_{\max} , cm^{-1}): 669 (C-S), 1116 (C=S), 1475 (Ar), 1514 (Ar), 1610 (Ar), 2875 (CH), 2936 (CH), 3410 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.96 (t, 3H, CH_3), 1.25-1.29 (m, 4H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 1.30-1.34 (m, 2H, CH_2CH_3 of hexyl group), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of hexyl group), 1.71-1.73 (m, 2H, naphthyl-O- CH_2CH_2), 1.94-1.96 (m, 2H, $\text{S-CH}_2\text{CH}_2$), 2.0 (bs, H, NH), 2.62-2.64 (m, 2H, NHCH_2), 2.82-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.02-4.06 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.95-7.62 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 375; Analysis: $\text{C}_{21}\text{H}_{29}\text{NOS}_2$, Calcd. (%): C, 67.15, H 7.78, N, 3.73; Obsd. (%): C, 67.59, H, 7.56, N, 3.51.

Octyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (10): Yield: 0.85 g (96.2 %); m.p.: 172 °C; IR (KBr, ν_{\max} , cm^{-1}): 667 (C-S), 1120 (C=S), 1475 (Ar), 1521 (Ar), 1612 (Ar), 2886 (CH), 2941 (CH), 3399 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.94 (t, 3H, CH_3), 1.27-1.29 (m, 8H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of octyl group), 1.32-1.34 (m, 2H, CH_2CH_3 of octyl group), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$ of *n*-octyl group), 2.0 (bs, H, NH), 2.62-2.64 (m, 2H, NHCH_2), 3.25-3.29 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.01-4.04 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.95-7.62 (m, 7H, Ar-H of naphthylloxy); Mass: m/e 375.17; Analysis: $\text{C}_{21}\text{H}_{29}\text{NOS}_2$, Calcd. (%): C, 67.15, H, 7.78, N, 3.73; O, 4.22; S, 17.04. Obsd. (%): C, 67.15, H, 7.78, N, 3.73, O, 4.26; S, 17.07.



Scheme-II

Octyl-dithiocarbamic acid-3-(naphthalene-2-yloxy)-propyl ester (11): Yield: 0.85 g (96.2 %); m.p.: 172 °C; IR (KBr, ν_{\max} , cm^{-1}): 667 (C-S), 1120 (C=S), 1472 (Ar), 1521 (Ar), 1612 (Ar), 2884 (CH), 2941 (CH), 3399 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.94 (t, 3H, CH_3), 1.27-1.29 (m, 8H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of octyl group), 1.30-1.32 (m, 2H, CH_2CH_3 of octyl group), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$ of *n*-octyl group), 2.0 (bs, H, NH), 2.38-2.42 (m, 2H, naphthyl-O- $\text{CH}_2\text{CH}_2\text{CH}_2\text{-S-C=S}$), 2.63-2.66 (m, 2H, NHCH_2), 2.83-2.87 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.01-4.04 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.98-7.66 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 389; Analysis: $\text{C}_{22}\text{H}_{31}\text{NOS}_2$, Calcd. (%): C, 67.62, H, 8.02, N, 3.59; Obsd. (%): C, 67.89, H, 7.90, N, 3.44.

Octyl-dithiocarbamic acid-4-(naphthalene-2-yloxy)-butyl ester (12): Yield: 0.86 g (94 %); m.p.: 156 °C; IR (KBr, ν_{\max} , cm^{-1}): 664 (C-S), 1108 (C=S), 1463 (Ar), 1514 (Ar), 1602 (Ar), 2862 (CH), 2926 (CH), 3392 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.93-0.95 (t, 3H, CH_3), 1.27-1.29 (m, 8H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of octyl group), 1.30-1.34 (m, 2H, CH_2CH_3 of octyl group), 1.52-1.57 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$), 2.0 (bs, H, NH), 2.63-2.67 (m, 2H, NHCH_2), 3.26-3.32 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.71-4.74 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 375; Analysis: $\text{C}_{21}\text{H}_{29}\text{NOS}_2$, Calcd. (%): C, 67.15, H 7.78, N, 3.73, Obsd. (%): C, 67.54, H, 7.56, N, 3.56.

Decyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (13): Yield: 0.86 g (94 %); m.p.: 156 °C; IR (KBr, ν_{\max} , cm^{-1}): 663 (C-S), 1108 (C=S), 1465 (Ar), 1511 (Ar), 1605 (Ar), 2862 (CH), 2926 (CH), 3392 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.94 (t, 3H, CH_3), 1.27-1.29 (m, 10H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of decyl group), 1.31-1.35 (m, 2H, CH_2CH_3 of decyl group), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$ of *n*-decyl group), 2.01 (bs, H, NH), 2.63-2.65 (m, 2H, NHCH_2), 3.25-3.29 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.97-7.64 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 403.25; Analysis: $\text{C}_{23}\text{H}_{33}\text{NOS}_2$, Calcd. (%): C, 68.40, H 8.20, N, 3.90, O, 3.92, S, 15.86; Obsd. (%): C, 68.44, H, 8.24, N, 3.96, O, 3.96, S, 15.89.

Decyl-dithiocarbamic acid-3-(naphthalene-2-yloxy)-propyl ester (14): Yield: 0.86 g (94 %); m.p.: 156 °C; IR (KBr, ν_{\max} , cm^{-1}): 664 (C-S), 1109 (C=S), 1462 (Ar), 1513 (Ar), 1602 (Ar), 2863 (CH), 2925 (CH), 3392 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.95 (t, 3H, CH_3), 1.27-1.29 (m, 10H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of decyl group), 1.31-1.35 (m, 2H, CH_2CH_3 of decyl group), 1.53-1.56 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$ of *n*-decyl group), 2.0 (bs, H, NH), 2.62-2.66 (m, 2H, NHCH_2), 2.84-2.87 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 2.34-2.38 (m, 2H, CH_2CH_2), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 417.67; Analysis: $\text{C}_{24}\text{H}_{35}\text{NOS}_2$, Calcd. (%): C, 69, H 8.40, N, 3.35, O, 3.80, S, 15.31. O, 3.83, S, 15.35. Obsd. (%): C, 69.02, H, 8.45, N, 3.35, O, 3.83, S, 15.35.

Decyl-dithiocarbamic acid-4-(naphthalene-2-yloxy)-butyl ester (15): Yield: 0.86 g (94 %); m.p.: 156 °C; IR (KBr, ν_{\max} , cm^{-1}): 663 (C-S), 1107 (C=S), 1462 (Ar), 1510 (Ar), 1606 (Ar), 2865 (CH), 2926 (CH), 3392 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 0.92-0.94 (t, 3H, CH_3), 1.27-1.29 (m, 10H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ of decyl group), 1.30-1.34 (m, 2H, CH_2CH_3 of decyl group), 1.52-1.58 (m, 2H, $\text{CH}_2\text{CH}_2\text{N}$ of *n*-decyl group), 2.0 (bs, H, NH), 2.62-2.66 (m, 2H, NHCH_2), 2.84-

2.87 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.93-1.95 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.67-1.71 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.97-7.64 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 431.70; Analysis: $\text{C}_{25}\text{H}_{37}\text{NOS}_2$, Calcd. (%): C, 69.52, H 8.63, N, 3.21, O, 3.70, S, 14.82. Obsd. (%): C, 69.56, H, 8.64, N, 3.24, O, 3.71, S, 14.86.

Pyrrolidine-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (16): Yield: 0.62 g (80.8 %); m.p.: 79 °C; IR (KBr, ν_{\max} , cm^{-1}): 657 (C-S), 1106 (C=S), 1454 (Ar), 1502 (Ar), 1600 (Ar), 2863 (CH), 2925 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.58-1.60 (m, 4H, CH_2 of pyrrolidine ring), 2.8 (t, 4H, CH_2N of pyrrolidine ring (2H, $\text{CH}_2\text{-S-C=S}$), 4.71-4.73 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 317; Analysis: $\text{C}_{17}\text{H}_{19}\text{NOS}_2$, Calcd. (%): C, 64.32, H, 6.03, N, 4.41, Obsd. (%): C, 63.85, H, 6.29, N, 4.64.

Pyrrolidine-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (17): Yield: 0.63 g (83.2 %); m.p.: 86 °C; IR (KBr, ν_{\max} , cm^{-1}): 672 (C-S), 1124 (C=S), 1475 (Ar), 1524 (Ar), 1605 (Ar), 2885 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.57-1.61 (m, 4H, CH_2 of pyrrolidine ring), 2.35-2.38 (m, 2H, naphthyl-O- $\text{CH}_2\text{CH}_2\text{CH}_2\text{-S-C=S}$), 2.8 (t, 4H, CH_2N of pyrrolidine ring), 2.82-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.02-4.04 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 331; Analysis: $\text{C}_{18}\text{H}_{21}\text{NOS}_2$, Calcd. (%): C, 65.22, H, 6.39, N, 4.23, Obsd. (%): C, 65.63, H, 6.12, N, 4.01.

Pyrrolidine-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (18): Yield: 0.64 g (86.5 %); m.p.: 95 °C; IR (KBr, ν_{\max} , cm^{-1}): 679 (C-S), 1125 (C=S), 1484 (Ar), 1525 (Ar), 1610 (Ar), 2885 (CH), 2947 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.56-1.60 (m, 4H, CH_2 of pyrrolidine ring), 1.71-1.72 (m, 2H, naphthyl-O- CH_2CH_2), 1.95-1.98 (m, 2H, $\text{S-CH}_2\text{CH}_2$), 2.8 (t, 4H, CH_2N of pyrrolidine ring), 2.84-2.88 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.02-4.05 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.97-7.64 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 345; Analysis: $\text{C}_{19}\text{H}_{23}\text{NOS}_2$, Calcd. (%): C, 66.09, H, 6.57, N, 4.15, Obsd. (%): C, 66.57, H, 6.32, N, 3.86.

Piperidine-1-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (19): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1110 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.48-1.50 (m, 6H, CH_2 of piperidine ring), 2.7 (t, 4H, CH_2N of piperidine ring), 3.28-3.30 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.71-4.73 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.95-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 331; Analysis: $\text{C}_{18}\text{H}_{21}\text{NOS}_2$, Calcd. (%): C, 65.22, H 6.39, N, 4.23, Obsd. (%): C, 65.73, H, 6.13, N, 3.98.

Piperidine-1-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (20): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1112 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2927 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.48-1.50 (m, 6H, CH_2 of piperidine ring), 2.7 (t, 4H, CH_2N of piperidine ring), 2.82-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 2.35-2.38 (m, 2H, CH_2CH_2), 4.01-4.04 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.95-7.62 (m, 7H, Ar-H of naphthyloxy); Mass: *m/e* 392.54; Analysis: $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_2\text{S}_3$, Calcd. (%): C, 58.22, H 6.20, N, 7.23, Obsd. (%): C, 58.14, H, 6.16, N, 7.14, O, 12.23, S, 16.13.

Piperidine-1-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (21): Yield: 0.66 g (82.5 %); m.p.: 89 °C;

IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1110 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2925 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 1.48-1.50 (m, 6H, CH_2 of piperidine ring), 2.7 (t, 4H, CH_2N of piperidine ring), 2.83-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.92-1.96 (m, 2H, CH_2CH_2), 1.68-1.71 (m, 2H, CH_2CH_2), 4.00-4.02 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.65 (m, 7H, Ar-H of naphthyl); Mass: m/e 406.56; Analysis: $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_3\text{S}_2$, Calcd. (%): C, 59.12, H 6.49, N, 6.82, O, 11.82, S, 15.75. Obsd. (%): C, 59.08, H, 6.45, N, 6.89, O, 11.89, S, 15.79.

4-Methyl-piperazine-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (22): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1112 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 2.24-2.27 (s, 3H, CH_3 of methyl piperazine ring), 2.44-2.48 (t, CH_2N of piperazine ring), 2.62-2.65 (t, CH_2N of piperazine ring), 2.0 (bs, H, NH), 3.25-3.29 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.65 (m, 7H, Ar-H of naphthyl); Mass: m/e 361; Analysis: $\text{C}_{18}\text{H}_{23}\text{N}_3\text{OS}_2$, Calcd. (%): C, 59.84, H 6.39, N, 11.60, O, 4.40, S, 17.72 % Obsd. (%): C, 59.80, H, 6.41, N, 11.62 % O, 4.43, S, 17.74.

4-Methyl-piperazine-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (23): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1110 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 2.24-2.27 (s, 3H, CH_3 of methyl piperazine ring), 2.44-2.48 (t, CH_2N of piperazine ring), 2.62-2.65 (t, CH_2N of piperazine ring), 2.0 (bs, H, NH), 2.84-2.87 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 2.34-2.38 (m, 2H, CH_2CH_2), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.63 (m, 7H, Ar-H of naphthyl); Mass: m/e 375.55; Analysis: $\text{C}_{19}\text{H}_{25}\text{N}_3\text{OS}_2$, Calcd. (%): C, 60.72, H 6.69, N, 11.23, O, 4.22, S, 17.05 % Obsd. (%): C, 60.76, H, 6.71, N, 11.19, O, 4.26, S, 17.08.

4-Methyl-piperazine-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (24): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1112 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2925 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 2.24-2.27 (s, 3H, CH_3 of methyl piperazine ring), 2.44-2.48 (t, CH_2N of piperazine ring), 2.62-2.65 (t, CH_2N of piperazine ring), 2.0 (bs, H, NH), 2.84-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.94-1.96 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.68-1.71 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.62 (m, 7H, Ar-H of naphthyl); Mass: m/e 389.58; Analysis: $\text{C}_{20}\text{H}_{27}\text{N}_3\text{OS}_2$, Calcd. (%): C, 61.64, H 6.93, N, 10.73, O, 4.08, S, 16.42 % Obsd. (%): C, 61.66, H, 6.99, N, 10.79, O, 4.11, S, 16.46.

Morpholine 4-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (25): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1110 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 3.62-3.67 (t, 2H, CH_2 of morpholine ring), 2.34-2.37 (t, CH_2N of morpholine ring), 2.35-2.40 (s, CH_2SCS), 2.84-2.86 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 2.34-2.38 (m, 2H, CH_2CH_2), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.95-7.62 (m, 7H, Ar-H of naphthyl); Mass: m/e 382.50; Analysis: $\text{C}_{17}\text{H}_{22}\text{N}_2\text{O}_4\text{S}_2$, Calcd. (%): C, 53.35, H 5.79, N, 7.30, O, 16.70, S, 16.72 % Obsd. (%): C, 53.38, H, 5.80, N, 7.32, O, 16.73, S, 16.77.

Morpholine 4-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (26): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1112 (C=S), 1461 (Ar), 1509

(Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 3.62-3.67 (t, 2H, CH_2 of morpholine ring), 2.34-2.37 (t, CH_2N of morpholine ring), 2.35-2.40 (s, $\text{CH}_2\text{S-C=S}$), 1.95-1.99 (m, 4H, CH_2CH_2), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.65 (m, 7H, Ar-H of naphthyl); Mass: m/e 396.52; Analysis: $\text{C}_{18}\text{H}_{24}\text{N}_2\text{O}_4\text{S}_2$, Calcd. (%): C, 45.50, H 6.09, N, 7.10, O, 16.10, S, 16.15. Obsd. (%): C, 45.52, H, 6.10, N, 7.06 % O, 16.14, S, 16.17.

Morpholine 4-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (27): Yield: 0.66 g (82.5 %); m.p.: 89 °C; IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1110 (C=S), 1461 (Ar), 1509 (Ar), 1605 (Ar), 2859 (CH), 2926 (CH); $^1\text{H NMR}$ (CDCl_3): δ = 3.62-3.67 (t, 2H, CH_2 of morpholine ring), 2.34-2.37 (t, CH_2N of morpholine ring), 2.35-2.40 (s, CH_2SCS), 3.25-3.28 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.94-1.96 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.68-1.71 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.96-7.63 (m, 7H, Ar-H of naphthyl); Mass: m/e 410.55; Analysis: $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_4\text{S}_2$, Calcd. (%): C, 55.55, H 6.39, N, 6.30, O, 15.55, S, 15.60. Obsd. (%): C, 55.58, H, 6.35, N, 6.85, O, 15.59, S, 15.62.

p-Tolyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (28): Yield: 0.76 g (88.4 %); m.p.: 137 °C; IR (KBr, ν_{\max} , cm^{-1}): 660 (C-S), 1111 (C=S), 1454 (Ar), 1502 (Ar), 1602 (Ar), 2851 (CH), 2928 (CH), 3388 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 2.34 (s, 3H, CH_3), 3.28-3.30 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.0 (bs, H, NH), 4.70-4.72 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.35-7.62 (m, 11H, Ar-H of naphthyl and phenyl ring); Mass: m/e 353; Analysis: $\text{C}_{20}\text{H}_{19}\text{NOS}_2$, Calcd. (%): C, 67.89, H, 5.45, N, 3.99. Obsd. (%): C, 67.63, H, 5.58, N, 4.12.

p-Tolyl-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (29): Yield: 0.76 g (88.4 %); m.p.: 137 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1111 (C=S), 1454 (Ar), 1502 (Ar), 1601 (Ar), 2851 (CH), 2928 (CH), 3388 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 2.34 (s, 3H, CH_3), 3.28-3.30 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.0 (bs, H, NH), 2.35-2.40 (s, $\text{CH}_2\text{-S-C=S}$), 1.95-1.99 (m, 4H, CH_2CH_2), 4.68-4.71 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.34-7.64 (m, 11H, Ar-H of naphthyl and phenyl ring); Mass: m/e 367.53; Analysis: $\text{C}_{21}\text{H}_{21}\text{NOS}_2$, Calcd. (%): C, 68.60, H, 5.72, N, 3.80, O, 4.31, S, 17.45 % Obsd. (%): C, 68.63, H, 5.76, N, 3.81 % O, 4.35, S, 17.45.

p-Tolyl-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (30): Yield: 0.76 g (88.4 %); m.p.: 137 °C; IR (KBr, ν_{\max} , cm^{-1}): 662 (C-S), 1111 (C=S), 1454 (Ar), 1502 (Ar), 1601 (Ar), 2851 (CH), 2928 (CH), 3388 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 2.34 (s, 3H, CH_3), 3.28-3.30 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 4.0 (bs, H, NH), 2.35-2.40 (s, CH_2SCS), 3.25-3.27 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 1.94-1.96 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.68-1.71 (m, 2H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 4.71-4.73 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$), 6.35-7.63 (m, 11H, Ar-H of naphthyl and phenyl ring); Mass: m/e 381.55; Analysis: $\text{C}_{22}\text{H}_{23}\text{NOS}_2$, Calcd. (%): C, 69.20, H, 6.06, N, 3.62, O, 4.15, S, 16.79. Obsd. (%): C, 69.25, H, 6.08, N, 3.67, O, 4.19, S, 16.81.

(4-Methoxy-4-phenyl)dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (31): Yield: 0.8 g (89.2 %); m.p.: 117 °C; IR (KBr, ν_{\max} , cm^{-1}): 659 (C-S), 1106 (C=S), 1455 (Ar), 1502 (Ar), 1600 (Ar), 2854 (CH), 2926 (CH), 3389 (NH); $^1\text{H NMR}$ (CDCl_3): δ = 3.28-3.30 (t, 2H, $\text{CH}_2\text{-S-C=S}$), 3.72 (s, 3H, OCH_3), 4.0 (bs, H, NH), 4.70-4.72 (t, 2H, $\text{CH}_2\text{-O-naphthyl}$),

6.35-7.64 (m, 11H, Ar-H of naphthoxy and phenyl ring); Mass: *m/e* 369; Analysis: C₂₀H₁₉NO₂S₂, Calcd. (%): C, 65.01, H, 5.18, N, 3.79, Obsd. (%): C, 65.47, H, 5.03, N, 3.48.

(4-Methoxy-4-phenyl)dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (32): Yield: 0.82 g (93.8 %); m.p.: 139 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1117 (C=S), 1472 (Ar), 1524 (Ar), 1614 (Ar), 2876 (CH), 2938 (CH), 3396 (NH); ¹H NMR (CDCl₃): δ = 2.38-2.42 (m, 2H, naphthyl-O-CH₂CH₂CH₂-S-C=S), 2.84-2.88 (t, 2H, CH₂-S-C=S), 3.74 (s, 3H, OCH₃), 4.0 (bs, H, NH), 4.02-4.05 (t, 2H, CH₂-O-naphthyl), 6.34-7.65 (m, 11H, Ar-H of naphthoxy and phenyl ring); Mass: *m/e* 383; Analysis: C₂₁H₂₁NO₂S₂, Calcd. (%): C, 65.76, H 5.52, N, 3.65, Obsd. (%): C, 65.27, H, 5.85, N, 3.81.

(4-Methoxy-4-phenyl)dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (33): Yield: 0.83 g (94.5 %); m.p.: 126 °C; IR (KBr, ν_{\max} , cm⁻¹): 681 (C-S), 1126 (C=S), 1484 (Ar), 1523 (Ar), 1610 (Ar), 2885 (CH), 2936 (CH), 3407 (NH); ¹H NMR (CDCl₃): δ = 1.71-1.74 (m, 2H, naphthyl-O-CH₂CH₂), 1.94-1.96 (m, 2H, S-CH₂CH₂), 2.01 (bs, H, NH), 2.82-2.86 (t, 2H, CH₂-S-C=S), 3.72 (s, 3H, OCH₃), 3.92-3.94 (d, 2H, CH₂ of benzylic proton), 4.01-4.04 (t, 2H, CH₂-O-naphthyl), 6.65-7.62 (m, 11H, Ar-H of naphthoxy and phenyl ring); Mass: *m/e* 411; Analysis: C₂₃H₂₅NO₂S₂, Calcd. (%): C, 67.12, H, 6.12, N, 3.40, Obsd. (%): C, 67.67, H, 6.40, N, 3.67.

Cyclohexyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (34): Yield: 0.714 g (85.5 %), m.p.: 112 °C; IR (KBr, ν_{\max} , cm⁻¹): 658 (C-S), 1103 (C=S), 1454 (Ar), 1502 (Ar), 1600 (Ar), 2851 (CH), 2926 (CH), 3373 (NH); ¹H NMR (CDCl₃): δ = 1.41-1.45 (m, 6H, CH₂ of cyclohexyl ring), 1.62-1.64 (m, 4H, CH₂ of cyclohexyl ring), 2.0 (bs, H, NH), 2.54-2.58 (m, H, tertiary H of cyclohexyl ring), 3.26-3.29 (t, 2H, CH₂-S-C=S), 4.71-4.74 (t, 2H, CH₂-O-naphthyl), 6.98-7.62 (m, 7H, Ar-H of naphthoxy); Mass: *m/e* 345; Analysis: C₁₉H₂₃NOS₂, Calcd. (%): C, 66.05, H, 6.71, N, 4.05, Obsd. (%): C, 65.65, H, 6.97, N, 4.23.

Cyclohexyl-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (35): Yield: 0.714 g (85.5 %); m.p.: 112 °C; IR (KBr, ν_{\max} , cm⁻¹): 658 (C-S), 1103 (C=S), 1454 (Ar), 1502 (Ar), 1600 (Ar), 2851 (CH), 2927 (CH), 3373 (NH); ¹H NMR (CDCl₃): δ = 1.44-1.48 (m, 6H, CH₂ of cyclohexyl ring), 1.63-1.66 (m, 4H, CH₂ of cyclohexyl ring), 2.0 (bs, H, NH), 2.54-2.59 (m, H, tertiary H of cyclohexyl ring), 3.28-3.30 (t, 2H, CH₂-S-C=S), 4.71-4.73 (t, 2H, CH₂-O-naphthyl), 6.95-7.62 (m, 7H, Ar-H of naphthoxy); Mass: *m/e* 345; Analysis: C₁₉H₂₃NOS₂, Calcd. (%): C, 66.05, H, 6.71, N, 4.05, Obsd. (%): C, 65.65, H, 6.97, N, 4.23.

Cyclohexyl-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (36): Yield: 0.75 g (94.5 %); m.p.: 126 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1121 (C=S), 1469 (Ar), 1523 (Ar), 1617 (Ar), 2879 (CH), 2937 (CH), 3408 (NH); ¹H NMR (CDCl₃): δ = 1.42-1.44 (m, 6H, CH₂ of cyclohexyl ring), 1.65-1.68 (m, 4H, CH₂ of cyclohexyl ring), 1.71-1.73 (m, 2H, naphthyl-O-CH₂CH₂), 1.94-1.96 (m, 2H, S-CH₂CH₂), 2.0 (bs, H, NH), 2.54-2.57 (m, H, tert. CH of cyclohexyl ring), 2.84-2.88 (t, 2H, CH₂-S-C=S), 4.02-4.06 (t, 2H, CH₂-O-naphthyl), 6.98-7.62 (m, 7H, Ar-H of naphthoxy); Mass: *m/e* 373; Analysis: C₂₁H₂₇NOS₂, Calcd. (%): C, 67.52, H, 7.28, N, 3.75, Obsd. (%): C, 67.84, H, 7.12, N, 3.59.

Benzyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)-ethyl ester (37): Yield: 0.75 g (87.3 %); m.p.: 101 °C; IR (KBr, ν_{\max} , cm⁻¹): 662 (C-S), 1112 (C=S), 1464 (Ar), 1512 (Ar), 1603 (Ar), 2865 (CH), 2926 (CH), 3385 (NH); ¹H NMR (CDCl₃): δ = 2.0 (bs, H, NH), 3.26-3.32 (t, 2H, CH₂-S-C=S), 3.92-3.94 (d, 2H, benzylic proton), 4.71-4.73 (t, 2H, CH₂-O-naphthyl), 6.98-7.65 (m, 12H, Ar-H of naphthoxy); Mass: *m/e* 353; Analysis: C₂₀H₁₉NOS₂, Calcd. (%): C, 67.95, H 5.42, N, 3.96, Obsd. (%): C, 67.63, H, 5.58, N, 4.12.

Benzyl-dithiocarbamic acid-3-(naphthalen-2-yloxy)-propyl ester (38): Yield: 0.75 g (89.8 %); m.p.: 109 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1114 (C=S), 1474 (Ar), 1514 (Ar), 1612 (Ar), 2863 (CH), 2926 (CH), 3398 (NH); ¹H NMR (CDCl₃): δ = 2.0 (bs, H, NH), 2.38-2.42 (m, 2H, naphthyl-O-CH₂CH₂CH₂-S-C=S), 2.82-2.86 (t, 2H, CH₂-S-C=S), 3.91-3.93 (d, 2H, CH₂ of benzylic hydrogens), 4.01-4.04 (t, 2H, CH₂-O-naphthyl), 6.96-7.62 (m, 12H, Ar-H of naphthoxy); Mass: *m/e* 367; Analysis: C₂₁H₂₁NOS₂, Calcd. (%): C, 68.63, H, 5.76, N, 3.81, Obsd. (%): C, 68.27, H, 5.94, N, 4.02.

Benzyl-dithiocarbamic acid-4-(naphthalen-2-yloxy)-butyl ester (39): Yield: 0.75 g (92.8 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm⁻¹): 673 (C-S), 1126 (C=S), 1484 (Ar), 1529 (Ar), 1612 (Ar), 2873 (CH), 2936 (CH), 3398 (NH); ¹H NMR (CDCl₃): δ = 1.71-1.74 (m, 2H, naphthyl-O-CH₂CH₂), 1.96-1.99 (m, 2H, S-CH₂CH₂), 2.0 (bs, H, NH), 2.84-2.88 (t, 2H, CH₂-S-C=S), 3.91-3.94 (d, 2H, CH₂ of benzylic proton), 4.01-4.04 (t, 2H, CH₂-O-naphthyl), 6.96-7.65 (m, 12H, Ar-H of naphthoxy); Mass: *m/e* 381; Analysis: C₂₂H₂₃NOS₂, Calcd. (%): C, 69.25, H 6.08, N, 3.67, Obsd. (%): C, 69.67, H, 5.87, N, 3.46.

Phenyl ethyl-dithiocarbamic acid-2-(naphthalen-2-yloxy)ethyl ester (40): Yield: 0.78 g (88.2 %); m.p.: 146 °C; IR (KBr, ν_{\max} , cm⁻¹): 661 (C-S), 1112 (C=S), 1464 (Ar), 1514 (Ar), 1605 (Ar), 2865 (CH), 2923 (CH), 3376 (NH); ¹H NMR (CDCl₃): δ = 2.0 (bs, H, NH), 2.80-2.82 (t, 2H, PhCH₂), 2.96-2.98 (m, 2H, NHCH₂CH₂Ph), 3.28-3.31 (t, 2H, CH₂-S-C=S), 4.71-4.73 (t, 2H, CH₂-O-naphthyl), 6.98-7.62 (m, 12H, Ar-H of naphthoxy and phenyl ring); Mass: *m/e* 367; Analysis: C₂₁H₂₁NOS₂, Calcd. (%): C, 68.63, H, 5.76, N, 3.81, Obsd. (%): C, 68.19, H, 6.06, N, 3.95.

Phenyl ethyl-dithiocarbamic acid-3-(naphthalen-2-yloxy)propyl ester (41): Yield: 0.8 g (91.4 %); m.p.: 172 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1126 (C=S), 1478 (Ar), 1519 (Ar), 1614 (Ar), 2878 (CH), 2933 (CH), 3396 (NH); ¹H NMR (CDCl₃): δ = 2.0 (bs, H, NH), 2.35-2.41 (m, 2H, naphthyl-O-CH₂CH₂CH₂-S-C=S), 2.80-2.82 (t, 2H, PhCH₂), 2.84-2.86 (t, 2H, CH₂-S-C=S), 2.96-3.02 (m, 2H, CH₂NH), 4.02-4.06 (t, 2H, CH₂-O-naphthyl), 6.95-7.62 (m, 12H, Ar-H of naphthoxy and phenyl group); Mass: *m/e* 381; Analysis: C₂₂H₂₃NOS₂, Calcd. (%): C, 69.25, H, 6.08, N, 3.67, Obsd. (%): C, 68.87, H, 6.29, N, 3.89.

Phenyl ethyl-dithiocarbamic acid-4-(naphthalen-2-yloxy)butyl ester (42): Yield: 0.8 g (94.8 %); m.p.: 179 °C; IR (KBr, ν_{\max} , cm⁻¹): 679 (C-S), 1149 (C=S), 1487 (Ar), 1533 (Ar), 1622 (Ar), 2884 (CH), 2944 (CH), 3438 (NH); ¹H NMR (CDCl₃): δ = 1.72-1.74 (m, 2H, naphthyl-O-CH₂CH₂), 1.96-1.97 (m, 2H, S-CH₂CH₂), 2.01 (bs, H, NH), 2.80-2.82 (t, 2H, PhCH₂), 2.86-2.88 (t, 2H, CH₂-S-C=S), 2.96-3.00 (m, 2H, CH₂NH), 4.02-4.06 (t, 2H, CH₂-O-naphthyl), 6.98-7.62 (m,

12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 395; Analysis: C₂₃H₂₅NOS₂, Calcd. (%): C, 69.93, H, 6.37, N, 3.54, Obsd. (%): C, 69.57, H, 6.55, N, 3.72.

Phenyl propyl-dithiocarbamic acid-2-(naphthalen-2-ylloxy)ethyl ester (43): Yield: 0.84 g (90.2 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1114 (C=S), 1462 (Ar), 1514 (Ar), 1600 (Ar), 2862 (CH), 2925 (CH), 3388 (NH); ¹H NMR (CDCl₃): δ = 1.86-1.88 (m, 2H, PhCH₂CH₂CH₂NH), 2.0 (bs, H, NH), 2.54-2.56 (t, 2H, PhCH₂), 2.65-2.64 (m, 2H, NHCH₂CH₂CH₂Ph), 3.28-3.31 (t, 2H, CH₂-S-C=S), 4.72-4.74 (t, 2H, CH₂-O-naphthyl), 6.95-7.62 (m, 12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 381; Analysis: C₂₂H₂₃NOS₂, Calcd. (%): C, 69.25, H 6.08, N, 3.67, Obsd. (%): C, 69.66, H, 5.99, N, 3.35.

Phenyl propyl-dithiocarbamic acid-3-(naphthalen-2-ylloxy)propyl ester (44): Yield: 0.84 g (93.2 %); m.p.: 135 °C; IR (KBr, ν_{\max} , cm⁻¹): 682 (C-S), 1129 (C=S), 1481 (Ar), 1533 (Ar), 1626 (Ar), 2884 (CH), 2936 (CH), 3416 (NH); ¹H NMR (CDCl₃): δ = 1.86-1.89 (m, 2H, Ph.CH₂CH₂CH₂), 2.0 (bs, H, NH), 2.38-2.42 (m, 2H, naphthyl-O-CH₂CH₂CH₂-S-C=S), 2.52-2.55 (t, 2H, PhCH₂), 2.62-2.64 (m, 2H, NHCH₂), 2.84-2.88 (t, 2H, CH₂-S-C=S), 4.02-4.05 (t, 2H, CH₂-O-naphthyl), 6.98-7.65 (m, 12H, Ar-H of naphthylloxy and phenyl group); Mass: *m/e* 395; Analysis: C₂₃H₂₅NOS₂, Calcd. (%): C, 69.83, H 6.37, N, 3.54, Obsd. (%): C, 69.34, H, 6.66, N, 3.74.

Phenyl propyl-dithiocarbamic acid-4-(naphthalen-2-ylloxy)butyl ester (45): Yield: 0.85 g (97.6 %); m.p.: 154 °C; IR (KBr, ν_{\max} , cm⁻¹): 692 (C-S), 1139 (C=S), 1486 (Ar), 1539 (Ar), 1628 (Ar), 2882 (CH), 2948 (CH), 3427 (NH); ¹H NMR (CDCl₃): δ = 1.71-1.73 (m, 2H, naphthyl-O-CH₂CH₂), 1.86-1.88 (m, 2H, PhCH₂CH₂CH₂NH), 1.96-1.99 (m, 2H, S-CH₂CH₂), 2.02 (bs, H, NH), 2.54-2.56 (t, 2H, PhCH₂), 2.64-2.68 (m, 2H, PhCH₂CH₂CH₂-N), 2.84-2.86 (t, 2H, CH₂-S-C=S), 2.98-3.01 (m, 2H, CH₂NH), 4.02-4.06 (t, 2H, CH₂-O-naphthyl), 6.98-7.62 (m, 12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 409; Analysis: C₂₄H₂₇NOS₂, Calcd. (%): C, 70.37, H, 6.64, N, 3.42, Obsd. (%): C, 69.95, H, 6.86, N, 3.62.

Di-*sec*-butyl-dithiocarbamic acid-2-(naphthalen-2-ylloxy)ethyl ester (46): Yield: 0.84 g (90.2 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1115 (C=S), 1462 (Ar), 1514 (Ar), 1601 (Ar), 2865 (CH), 2926 (CH), 3388 (NH); ¹H NMR (CDCl₃): δ = 1.08-1.10 (d, CH₃), 2.75-2.79 (m, 8H, CH₃CH₂CH₃), 1.38-1.41 (M, 6H, CH₃CHCH₃), 0.94-0.96 (t, 2H, CH₃CHCH₂), 3.28-3.30 (t, 2H, CH₂-S-C=S), 4.71-4.73 (t, 2H, CH₂-O-naphthyl), 6.95-7.63 (m, 12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 375.59; Analysis: C₂₁H₂₉NOS₂, Calcd. (%): C, 67.10, H 7.75, N, 3.70, O, 4.22, S, 17.05. Obsd. (%): C, 67.15, H, 7.78, N, 3.73, O, 4.26, S, 17.07.

Di-*sec*-butyl-dithiocarbamic acid-3-(naphthalen-2-ylloxy)propyl ester (47): Yield: 0.84 g (90.2 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1114 (C=S), 1462 (Ar), 1514 (Ar), 1600 (Ar), 2862 (CH), 2925 (CH), 3388 (NH); ¹H NMR (CDCl₃): δ = 1.08-1.10 (d, CH₃), 2.75-2.79 (m, 8H, CH₃CH₂CH₃), 1.38-1.41 (M, 6H, CH₃CHCH₃), 0.94-0.96 (t, 2H, CH₃CHCH₂), 3.28-3.30 (t, 2H, CH₂-S-C=S), 2.35-2.39 (m, 2H, CH₂CH₂CH₂), 4.72-4.74 (t, 2H, CH₂-O-naphthyl), 6.96-7.62 (m, 12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 389.62; Analysis: C₂₂H₃₁NOS₂, Calcd. (%): C, 67.80, H

8.01, N, 3.60, O, 4.10, S, 16.42. Obsd. (%): C, 67.82, H, 8.02, N, 3.59, O, 4.11, S, 16.46.

Di-*sec*-butyl-dithiocarbamic acid-4-(naphthalen-2-ylloxy)butyl ester (48): Yield: 0.84 g (90.2 %); m.p.: 119 °C; IR (KBr, ν_{\max} , cm⁻¹): 668 (C-S), 1114 (C=S), 1465 (Ar), 1514 (Ar), 1600 (Ar), 2865 (CH), 2927 (CH), 3388 (NH); ¹H NMR (CDCl₃): δ = 1.08-1.10 (d, CH₃), 2.75-2.79 (m, 8H, CH₃CH₂CH₃), 1.38-1.41 (M, 6H, CH₃CHCH₃), 0.94-0.96 (t, 2H, CH₃CHCH₂), 3.27-3.31 (t, 2H, CH₂-S-C=S), 1.92-1.96 (m, 2H, CH₂CH₂CH₂), 1.68-1.71 (m, 2H, CH₂CH₂CH₂), 4.71-4.73 (t, 2H, CH₂-O-naphthyl), 6.97-7.64 (m, 12H, Ar-H of naphthylloxy and phenyl ring); Mass: *m/e* 403.64; Analysis: C₂₃H₃₃NOS₂, Calcd. (%): C, 68.41, H 8.20, N, 3.44, O, 3.92, S, 15.87. Obsd. (%): C, 68.44, H, 8.24, N, 3.47, O, 3.96, S, 15.89.

RESULTS AND DISCUSSION

The synthetic route of ω -substituted 2-naphthylloxy haloalkanes and desired products (**4-48**) as shown in **Scheme-I** is prepared by direct condensation of β -naphthol and alkyl dihalide. Intermediate ω -substituted 2-naphthylloxy haloalkanes (**3**) was prepared by reacting alkyl dihalide (**2**) with β -naphthol (**1**) in the presence of anhydrous K₂CO₃ which was subsequently converted to corresponding dithiocarbamates of **4** to **48** by reaction of various types of primary and secondary amines involving Triton B/CS₂ system at room temperature. Hence, dithiocarbamates of desired (**4-48**) series is synthesized employing various kinds of aliphatic, aromatic, alicyclic, heterocyclic primary and secondary amines. The protocol proved to be successful and the desired product was isolated and further confirmed by spectroscopic and analytical methods.

The structural characterization of title compounds have been done by melting point, ¹H NMR, ¹³C NMR and high-resolution mass spectrometry (HRMS). All spectral data were consistent with the assigned structures. The comparative study of the yield of **3a**, **3b** and **3c** increases with number of carbon due to +I effect. However, final yield of dithiocarbamates of Prototype **I** (**4-48**) is dependent upon the electron releasing effect of the amines like pyrrolidine, piperidine, *N*-methyl piperazine, cyclohexane, phenyl ethyl amine and phenyl propyl amine show higher yield compared to primary amines (Table-1).

Antimicrobial screening: The series of compounds were screened for antimicrobial activity through microdilution method using various bacterial and fungal strains. The antifungal and antibacterial values were estimated as MIC values. Fluconazole and ciprofloxacin were used as the standard antifungal and antibacterial drug. As shown in Table-2, the SAR of these compounds can be studied by varying the alkyl chain and amines attached to these range of compounds. Compounds having three-carbon chain attached to them are found to be more active as compared to two-carbon or four-carbon chain. Among **28**, **29**, **30** and **31**, **32**, **33**, **29** and **33** were found to possess higher potency as compared to others because of the three-carbon chain attached to them. The higher potency of three-carbon chain is due to hydrophilicity. Upon studying the effect of various types of amines, we found that compounds like **28**, **29**, **30**, **31**, **32** and **33** having aromatic amine like anisidine and toluidine possessed comparable values to control drugs. Substitution of heterocyclic amines in compounds **16**,

TABLE-1
SYNTHESIS OF VARIOUS TYPES OF DITHIOCARBAMATES OF ω -SUBSTITUTED 2-NAPHTHYLOXY ALKANES

Comp. No.	n	R ₁	R ₂	Time (min)	Yield (%)
4	1	C ₄ H ₉	H	35	93
5	2	C ₄ H ₉	H	30	95
6	3	C ₄ H ₉	H	38	93
7	1	C ₆ H ₁₁	H	30	94
8	2	C ₆ H ₁₁	H	30	95
9	3	C ₆ H ₁₁	H	35	94
10	1	C ₈ H ₁₅	H	40	90
11	2	C ₈ H ₁₅	H	35	94
12	3	C ₈ H ₁₅	H	38	92
13	1	C ₁₀ H ₁₉	H	40	93
14	2	C ₁₀ H ₁₉	H	38	95
15	3	C ₁₀ H ₁₉	H	40	92
16	1	R ₁ = R ₂ = Pyrrolidine	–	25	96
17	2	R ₁ = R ₂ = Pyrrolidine	–	20	98
18	3	R ₁ = R ₂ = Pyrrolidine	–	25	95
19	1	R ₁ = R ₂ = Piperidine	–	30	98
20	2	R ₁ = R ₂ = Piperidine	–	28	95
21	3	R ₁ = R ₂ = Piperidine	–	30	98
22	1	R ₁ = R ₂ = <i>N</i> -methyl piperazine	–	20	98
23	2	R ₁ = R ₂ = <i>N</i> -methyl piperazine	–	20	98
24	3	R ₁ = R ₂ = <i>N</i> -methyl piperazine	–	20	96
25	1	R ₁ = R ₂ = Morpholine	–	25	90
26	2	R ₁ = R ₂ = Morpholine	–	20	92
27	3	R ₁ = R ₂ = Morpholine	–	25	90
28	1	R ₁ = R ₂ = Toludine	–	25	92
29	2	R ₁ = R ₂ = Toludine	–	20	94
30	3	R ₁ = R ₂ = Toludine	–	25	90
31	1	R ₁ = R ₂ = Anisidine	–	30	92
32	2	R ₁ = R ₂ = Anisidine	–	20	90
33	3	R ₁ = R ₂ = Anisidine	–	25	92
34	1	R ₁ = R ₂ = Cyclohexane	–	25	95
35	2	R ₁ = R ₂ = Cyclohexane	–	25	92
36	3	R ₁ = R ₂ = Cyclohexane	–	30	94
37	1	Ph(CH ₂)	H	40	93
38	2	Ph(CH ₂)	H	25	96
39	3	Ph(CH ₂)	H	30	98
40	1	Ph(CH ₂ CH ₂)	H	20	98
41	2	Ph(CH ₂ CH ₂)	H	20	98
42	3	Ph(CH ₂ CH ₂)	H	25	90
43	1	Ph(CH ₂ CH ₂ CH ₂)	H	25	92
44	2	Ph(CH ₂ CH ₂ CH ₂)	H	30	92
45	3	Ph(CH ₂ CH ₂ CH ₂)	H	25	95
46	1	R ₁ = R ₂ = Dibutyl	–	30	97
47	2	R ₁ = R ₂ = Dibutyl	–	25	95
48	3	R ₁ = R ₂ = Dibutyl	–	35	80

17, 18, 19, 20, 21, 22, 23, 24, 25, 26 and 27 gave promising result. Among substituted aromatic amines benzyl amine (37, 38 and 39) gave better result than phenyl ethyl (40, 41 and 42) and phenyl propyl amine (43, 44 and 45).

Conclusion

In conclusion, a convenient and efficient protocol for one-pot synthesis has been developed, employing three components coupling of various amines with variety of *via* CS₂ Bridge using Triton-B. This method produces the corresponding dithiocarbamates in good to excellent yields. Furthermore, the compounds produced by this method exhibited maximum potency for antifungal and antibacterial activity which is comparable to standard drug.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge fruitful discussions with Dr. Devdutt Chaturvedi and Amity University Uttar Pradesh (AUUP), Lucknow Campus, Lucknow, U.P., for their constant encouragement and support for research. Financial support from the Department of Science and Technology (DST), Govt. of India (Grant No.SR/FT/CS-147/2010) is gratefully acknowledged.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interests regarding the publication of this article.

TABLE-2
ANTIMICROBIAL DATA OF DITHIOCARBAMATES OF ω -SUBSTITUTED 2-NAPHTHYLOXY ALKANES

Compounds	Antifungal activity				Antibacterial activity			
	<i>C. albicans</i>	<i>S. schenckii</i>	<i>A. fumigatus</i>	<i>A. niger</i>	<i>S. aureus</i>	MRSA*	<i>E. coli</i>	<i>P. aeruginosa</i>
4	1.56	1.52	6.25	8	6.25	>50	3.12	6.25
5	1.58	1.56	6.27	8.5	6.27	>50	3.14	6.26
6	1.60	1.58	6.28	9	6.28	>50	3.18	6.28
7	3.12	1.56	>50	8	8	>50	6.25	>50
8	3.14	1.58	>50	8.5	8.8	>50	6.27	>50
9	3.18	1.62	>50	8.8	9	>50	6.28	>50
10	6.25	6.25	>50	16	16	>50	>50	>50
11	6.27	6.27	>50	15	17	>50	>50	>50
12	6.30	6.28	>50	18	18	>50	>50	>50
13	12.5	6.25	>50	>50	>50	>50	>50	>50
14	12.8	6.28	>50	>50	>50	>50	>50	>50
15	13.0	6.28	>50	>50	>50	>50	>50	>50
16	0.92	0.32	0.80	1.30	3.12	>50	0.62	3.12
17	0.75	0.28	0.71	1.20	3.14	>50	0.48	3.14
18	0.82	0.32	0.78	1.25	3.18	>50	0.54	3.16
19	0.95	0.35	0.89	25	6.25	>50	0.84	6.25
20	0.80	0.30	0.78	24	6.27	>50	0.55	6.26
21	0.88	0.33	0.85	28	6.28	>50	0.72	6.28
22	0.88	0.30	0.75	0.82	0.78	50	0.42	1.38
23	0.70	0.26	0.66	0.64	0.64	50	0.24	1.33
24	0.80	0.28	0.70	0.74	0.70	50	0.36	1.35
25	0.81	0.29	0.68	0.66	0.76	30	0.32	1.25
26	0.65	0.20	0.59	0.52	0.60	24	0.20	1.27
27	0.74	0.25	0.62	0.58	0.68	28	0.25	1.29
28	0.80	0.25	0.62	0.58	0.68	27	0.30	0.72
29	0.64	0.17	0.55	0.45	0.57	20	0.15	0.52
30	0.72	0.20	0.58	0.50	0.61	25	0.19	0.65
31	0.78	0.20	0.52	0.50	0.59	22	0.20	0.68
32	0.50	0.15	0.45	0.40	0.45	18	0.10	0.48
33	0.65	0.18	0.50	0.45	0.55	20	0.15	0.55
34	2.89	3.12	>50	>50	8	>50	8	12.5
35	2.90	3.14	>50	>50	8.5	>50	8.5	12.7
36	3.28	3.20	>50	>50	9	>50	9	13.0
37	1.05	>50	50	>50	>50	>50	50	8
38	0.95	>50	50	>50	>50	>50	50	8.8
39	1.10	>50	50	>50	>50	>50	50	9
40	1.16	>50	>50	>50	>50	>50	>50	>50
41	1.05	>50	>50	>50	>50	>50	>50	>50
42	1.22	>50	>50	>50	>50	>50	>50	>50
43	>50	>50	>50	>50	>50	>50	>50	>50
44	>50	>50	>50	>50	>50	>50	>50	>50
45	>50	>50	>50	>50	>50	>50	>50	>50
46	>50	>50	>50	>50	>50	>50	>50	>50
47	>50	>50	>50	>50	>50	>50	>50	>50
48	>50	>50	>50	>50	>50	>50	>50	>50
Fluconazole	0.25	0.09	0.35	0.25	–	–	–	–
Ciprofloxacin	–	–	–	–	0.25	16	0.03	0.25

REFERENCES

- S.J. Shaw, *Mini Rev. Med. Chem.*, **8**, 276 (2008); <https://doi.org/10.2174/138955708783744137>.
- D.K. Hutchinson, *Curr. Top. Med. Chem.*, **3**, 1021 (2003); <https://doi.org/10.2174/1568026033452195>.
- T. Asaka, A. Manaka and H. Sugiyama, *Curr. Top. Med. Chem.*, **3**, 961 (2003); <https://doi.org/10.2174/1568026033452140>.
- D. Sze, K. Miller and B. Neilan, *Recent Patents Anticancer. Drug Discov.*, **3**, 14 (2008); <https://doi.org/10.2174/157489208783478685>.
- D.J. Halls, *Mikrochim. Acta*, **57**, 62 (1969); <https://doi.org/10.1007/BF01216666>.
- T.W. Greene and T.P.G.M. Wuts, *Protective Group in Organic Synthesis*. John Wiley & Sons, Inc.: New York, edn 4 (2007);
- P.J. Kocienski, *Protective Groups*, Thieme-Verlag, Stuttgart, edn 3 (2003).
- J.P. Mayer, G.S. Lewis, M.J. Curtis and J. Zhang, *Tetrahedron Lett.*, **38**, 8445 (1997); [https://doi.org/10.1016/S0040-4039\(97\)10276-3](https://doi.org/10.1016/S0040-4039(97)10276-3).
- H.P. Buchstaller, *Tetrahedron*, **54**, 3465 (1998); [https://doi.org/10.1016/S0040-4020\(98\)00079-9](https://doi.org/10.1016/S0040-4020(98)00079-9).
- P.T. Holte, L. Thijs and B. Zwanenburg, *Tetrahedron Lett.*, **39**, 7404 (1998); [https://doi.org/10.1016/S0040-4039\(98\)01608-6](https://doi.org/10.1016/S0040-4039(98)01608-6).
- Y. Liu, K. Cui, W. Lu, W. Luo, J. Wang, J. Huang and C. Guo, *Molecules*, **16**, 4527 (2011); <https://doi.org/10.3390/molecules16064527>.

12. F. Ozkanli, A.G. Ozkanli, K. Ozadali, E. Yildirim and K. Erol, *FABAD J. Pharm. Sci.*, **35**, 19 (2010).
13. L. Yurttas, Y. Özkay, M. Duran, G. Turan-Zitouni, A. Özdemir, Z. Cantürk, K. Küçükoglu and Z.A. Kaplancikli, *Phosphorus Sulphur Silicon Rel. Elem.*, **191**, 1166 (2016); <https://doi.org/10.1080/10426507.2016.1150277>.
14. K.M. Heda and S.P. Deshmukh, *Int. J. Chem. Sci.*, **9**, 1705 (2011).
15. K. Bacharaju, S.R. Jambula, S. Sivan, S.J. Tangeda and V. Manga, *Bioorg. Med. Chem. Lett.*, **22**, 3274 (2012); <https://doi.org/10.1016/j.bmcl.2012.03.018>.
16. Y. Horita, T. Takii, R. Kuroishi, T. Chiba, K. Ogawa, L. Kremer, Y. Sato, Y.S. Lee, T. Hasegawa and K. Onozaki, *Bioorg. Med. Chem.*, **21**, 899 (2011); <https://doi.org/10.1016/j.bmcl.2010.12.084>.
17. Y. Zou, S. Yu, R. Li, Q. Zhao, X. Li, M. Wu, T. Huang, X. Chai, H. Hu and Q. Wu, *Eur. J. Med. Chem.*, **74**, 366 (2014); <https://doi.org/10.1016/j.ejmech.2014.01.009>.
18. S.L. Cao, Y. Wang, L. Zhu, J. Liao, Y.W. Guo, L.L. Chen, H.Q. Liu and X. Xu, *Eur. J. Med. Chem.*, **45**, 3850 (2010); <https://doi.org/10.1016/j.ejmech.2010.05.038>.
19. M.A.-H. Zahran, T.A.-R. Salem, R.M. Samaka, H.S. Agwa and A.R. Awad, *Bioorg. Med. Chem.*, **16**, 9708 (2008); <https://doi.org/10.1016/j.bmc.2008.09.071>.
20. J. Coro, R. Atherton, S. Little, H. Wharton, V. Yardley, A. Alvarez Jr., M. Suarez, R. Perez and H. Rodriguez, *Bioorg. Med. Chem.*, **16**, 1312 (2006); <https://doi.org/10.1016/j.bmcl.2005.11.060>.
21. V. Marakov, O.B. Riabova, A. Yuschenko, N. Urlyapova, A. Daudova, P.F. Ziplef and U. Mollmann, *J. Antimicrob. Chemother.*, **57**, 1134 (2006); <https://doi.org/10.1093/jac/dkl095>.
22. S.L. Cao, Y.P. Feng, X.L. Zheng, Y.Y. Jiang, M. Zhang, Y. Wang and M. Xu, *Arch. Pharm. Chem. Life Sci.*, **339**, 250 (2006); <https://doi.org/10.1002/ardp.200500264>.
23. X. Hou, Z. Ge, T. Wang, W. Guo, J. Wu, J. Cui, C. Lai and R. Li, *Arch. Pharm. Chem. Life Sci.*, **344**, 320 (2011); <https://doi.org/10.1002/ardp.201000259>.
24. A. Goel, S.J. Mazur, R.J. Fattah, T.L. Hartman, J.A. Turpin, M. Huang, W.G. Rice, E. Appella and J.K. Inman, *Bioorg. Med. Chem. Lett.*, **12**, 767 (2002); [https://doi.org/10.1016/S0960-894X\(02\)00007-0](https://doi.org/10.1016/S0960-894X(02)00007-0).
25. X.Y. He, L. Lu, J. Qiu, P. Zou, F. Yu, X.K. Jiang, L. Li, S. Jiang, S. Liu and L. Xie, *Bioorg. Med. Chem.*, **21**, 7539 (2013); <https://doi.org/10.1016/j.bmc.2013.04.046>.
26. S.L. Cao, Y. Han, C.Z. Yuan, Y. Wang, Z.K. Xiahou, J. Liao, R.T. Gao, B.B. Mao, B.L. Zhao, Z.F. Li and X. Xu, *Eur. J. Med. Chem.*, **64**, 401 (2013); <https://doi.org/10.1016/j.ejmech.2013.04.017>.
27. S. Jangir, V. Bala, N. Lal, L. Kumar, A. Sarswat, A. Kumar, Hamidullah, K.S. Saini, V. Sharma, V. Verma, J.P. Maikhuri, R. Konwar, G. Gupta and V.L. Sharma, *Eur. J. Med. Chem.*, **85**, 638 (2014); <https://doi.org/10.1016/j.ejmech.2014.08.028>.
28. P. Liu, C. Li, J. Zhang and X. Xu, *Synth. Commun.*, **43**, 3342 (2013); <https://doi.org/10.1080/00397911.2013.783600>.
29. A.Z. Halimehjani, Y. Pourshojaei and M.R. Saidi, *Tetrahedron Lett.*, **50**, 32 (2009); <https://doi.org/10.1016/j.tetlet.2008.10.063>.
30. J. Jamir, U.B. Sinha, J. Nath and B.K. Patel, *Synth. Commun.*, **42**, 951 (2012); <https://doi.org/10.1080/00397911.2010.532276>.
31. M. Kienle, A. Unsinn and P. Knochel, *Angew. Chem. Int. Ed.*, **49**, 4751 (2010); <https://doi.org/10.1002/anie.201001025>.
32. V. Aucagne, C. Lorin, A. Tatibouet and P. Rollin, *Tetrahedron Lett.*, **46**, 4349 (2005); <https://doi.org/10.1016/j.tetlet.2005.04.112>.
33. K. Alagiri and K.R. Prabhu, *Chem. Eur. J.*, **17**, 6922 (2011); <https://doi.org/10.1002/chem.201100817>.
34. N.K. Kumar, K. Sreeramamurthy, S. Palle, K. Muktanti and P. Das, *Tetrahedron Lett.*, **51**, 899 (2010); <https://doi.org/10.1016/j.tetlet.2009.11.127>.
35. M.A. Khalilzadeh, Z. Hossaini, M.M. Baradarani and A. Hasannia, *Tetrahedron Lett.*, **66**, 8464 (2010); <https://doi.org/10.1016/j.tet.2010.08.041>.
36. J.Y. Park, I.A. Ryu, H.H. Park, D.C. Ha and Y.D. Gong, *Synthesis*, 913 (2009); <https://doi.org/10.1055/s-0028-1087970>.
37. D. Katiyar, V.K. Tiwari, R.P. Tripathi, A. Srivastava, V. Chaturvedi, R. Srivastava and B.S. Srivastava, *Bioorg. Med. Chem.*, **11**, 4369 (2003); [https://doi.org/10.1016/S0968-0896\(03\)00480-2](https://doi.org/10.1016/S0968-0896(03)00480-2).
38. P. Das, C.K. Kumar, N. Kumar, M. Innus, J. Iqbal and N. Srinivas, *Tetrahedron Lett.*, **49**, 992 (2008);
39. U.A. Mohsen, *Cukurova Med. J.*, **39**, 729 (2014); <https://doi.org/10.17826/cutf.79473>.
40. S.K. Tandel, S. Rajappa and S.V. Pansare, *Tetrahedron*, **49**, 7479 (1993); [https://doi.org/10.1016/S0040-4020\(01\)87224-0](https://doi.org/10.1016/S0040-4020(01)87224-0).
41. F.R. Charati, Z. Hossaini and R. Hajinasiri, *J. Appl. Chem. Res.*, **20**, 54 (2012).
42. L.J. Bahrain, P.J. Jones and H. Hopf, *Beilstein J. Org. Chem.*, **8**, 1999 (2012); <https://doi.org/10.3762/bjoc.8.226>.
43. N.S. Gu'nay, G. Capan, N. Ulusoy, N. Ergenc, G. Ötük and D. Kaya, *Il Farmaco*, **54**, 826 (1999); [https://doi.org/10.1016/S0014-827X\(99\)00109-3](https://doi.org/10.1016/S0014-827X(99)00109-3).
44. K. Chauhan, M. Sharma, P. Singh, V. Kumar, P.K. Shukla, M.I. Siddiqi and P.M.S. Chauhan, *MedChemComm*, **3**, 1104 (2012); <https://doi.org/10.1039/c2md20109g>.
45. P.M. Madalageri and O. Kotresh, *Der Pharm. Chem.*, **4**, 2187 (2012).
46. C. Donnici, L. Nogueira, M. Araujo, S. Oliveira, T. Magalhães, M. Lopes, A. Silva, A. Ferreira, C. Martins and M. de Resende Stoianoff, *Molecules*, **19**, 5402 (2014); <https://doi.org/10.3390/molecules19045402>.
47. E. Septolu, M.D. Aytém, E. Killic, M. Ozalp and U. Califi, *FABAD J. Pharm. Sci.*, **29**, 1 (2004).
48. D. Chaturvedi, S. Ray, A.K. Srivastava and R. Chander, *Bioorg. Med. Chem.*, **16**, 2489 (2008); <https://doi.org/10.1016/j.bmc.2007.11.062>.